

How to obtain lower bounds in set theory

Sandra Müller

Universität Wien

May 8, 2020

CUNY Set Theory Seminar



Inner Model Theory

Determinacy

Combinatorics of κ -
Trees

Looking beyond

The main goal of inner model theory is to construct L -like models, which we call mice, for stronger and stronger large cardinals.

Definition

Let E be a set or a proper class. Let

$$\begin{aligned}J_0[E] &= \emptyset \\J_{\alpha+\omega}[E] &= \text{rud}_E(J_\alpha[E] \cup \{J_\alpha[E]\}) \\J_{\omega\lambda}[E] &= \bigcup_{\alpha < \lambda} J_{\omega\alpha}[E] \text{ for limit } \lambda \\L[E] &= \bigcup_{\alpha \in \text{Ord}} J_{\omega\alpha}[E]\end{aligned}$$

Note that rud_E denotes the closure under functions which are rudimentary in E (i.e. basic set operations like minus, union and pairing or intersection with E).

Condensation Let α be an infinite limit ordinal and let

$$M \prec (J_\alpha, \in).$$

Then the transitive collapse of M is equal to J_β for some limit ordinal $\beta \leq \alpha$.

Condensation Let α be an infinite limit ordinal and let

$$M \prec (J_\alpha, \in).$$

Then the transitive collapse of M is equal to J_β for some limit ordinal $\beta \leq \alpha$.

Comparison Let J_α and J_β for limit ordinals α and β be initial segments of L . Then one is an initial segment of the other, that means

$$J_\alpha \trianglelefteq J_\beta \text{ or } J_\beta \trianglelefteq J_\alpha.$$

A First Equivalence for Analytic Determinacy

Definition

Let x be a real. We say $x^\#$ exists iff for some limit ordinal λ , the model $J_{\omega\lambda}[x]$ has an uncountable set Γ of indiscernibles, i.e. for $n < \omega$ and any two increasing sequences $(\alpha_0, \dots, \alpha_n)$ and $(\beta_0, \dots, \beta_n)$ from Γ and any formula φ ,

$$J_{\omega\lambda}[x] \models \varphi(x, \alpha_0, \dots, \alpha_n) \Leftrightarrow J_{\omega\lambda}[x] \models \varphi(x, \beta_0, \dots, \beta_n).$$

A First Equivalence for Analytic Determinacy

Definition

Let x be a real. We say $x^\#$ exists iff for some limit ordinal λ , the model $J_{\omega\lambda}[x]$ has an uncountable set Γ of indiscernibles, i.e. for $n < \omega$ and any two increasing sequences $(\alpha_0, \dots, \alpha_n)$ and $(\beta_0, \dots, \beta_n)$ from Γ and any formula φ ,

$$J_{\omega\lambda}[x] \models \varphi(x, \alpha_0, \dots, \alpha_n) \Leftrightarrow J_{\omega\lambda}[x] \models \varphi(x, \beta_0, \dots, \beta_n).$$

Theorem (Harrington, Martin)

The following are equivalent.

- (a) *All analytic games are determined.*
- (b) *$x^\#$ exists for all reals x .*

A First Equivalence for Analytic Determinacy

Definition

Let x be a real. We say $x^\#$ exists iff for some limit ordinal λ , the model $J_{\omega\lambda}[x]$ has an uncountable set Γ of indiscernibles, i.e. for $n < \omega$ and any two increasing sequences $(\alpha_0, \dots, \alpha_n)$ and $(\beta_0, \dots, \beta_n)$ from Γ and any formula φ ,

$$J_{\omega\lambda}[x] \models \varphi(x, \alpha_0, \dots, \alpha_n) \Leftrightarrow J_{\omega\lambda}[x] \models \varphi(x, \beta_0, \dots, \beta_n).$$

Theorem (Harrington, Martin)

The following are equivalent.

- (a) *All analytic games are determined.*
- (b) *$x^\#$ exists for all reals x .*

To see how this relates to measurable cardinals, we need to look at a different definition of $x^\#$.

Definition (Equivalent definition of $x^\#$)

If it exists, let $x^\#$ be the unique premouse of the form $\mathcal{M} = (J_\alpha[x], \in, U)$ with $\text{crit}(U) = \kappa$ such that

- 1 if $z \subset \mathcal{P}(\kappa)$ is in \mathcal{M} with $|z|^\mathcal{M} = \kappa$, then $U \cap z \in \mathcal{M}$,
- 2 $\mathcal{M} \models U$ is a non-trivial normal κ -complete ultrafilter on κ ,
- 3 a fine structural condition which implies that $(J_{\alpha+\omega}[x], \in, U) \models |\alpha| = \omega$, and
- 4 every countable linear iterate of \mathcal{M} (via U) is well-founded.

Definition (Equivalent definition of $x^\#$)

If it exists, let $x^\#$ be the unique premouse of the form $\mathcal{M} = (J_\alpha[x], \in, U)$ with $\text{crit}(U) = \kappa$ such that

- 1 if $z \subset \mathcal{P}(\kappa)$ is in \mathcal{M} with $|z|^{\mathcal{M}} = \kappa$, then $U \cap z \in \mathcal{M}$,
- 2 $\mathcal{M} \models U$ is a non-trivial normal κ -complete ultrafilter on κ ,
- 3 a fine structural condition which implies that $(J_{\alpha+\omega}[x], \in, U) \models |\alpha| = \omega$, and
- 4 every countable linear iterate of \mathcal{M} (via U) is well-founded.

Note: The images of the critical point κ of the external measure U (when iterating the model $x^\#$ linearly) form an uncountable set of indiscernibles Γ for some large enough $J_{\omega\lambda}[x]$.

Basic Concepts of Inner Model Theory

To generalize this to larger large cardinals we need to study the underlying concepts a bit more.

Basic Concepts of Inner Model Theory

To generalize this to larger large cardinals we need to study the underlying concepts a bit more.

Mitchell and Jensen generalized the concept of measures to extenders to obtain stronger ultrapowers.

Definition

Let \mathcal{M} be a countable model of set theory. An *extender* over \mathcal{M} is a system of ultrafilters whose ultrapowers form a directed system, such that they give rise to a single elementary embedding.

In fact for every embedding $j : \mathcal{M} \rightarrow \mathcal{N}$ there is an extender E over \mathcal{M} which gives rise to this embedding.

Comparison

One key concept of inner model theory is building *iterated ultrapowers* to compare two models.

Comparison


One key concept of inner model theory is building *iterated ultrapowers* to compare two models.



\mathcal{M}

Comparison

One key concept of inner model theory is building *iterated ultrapowers* to compare two models.




Comparison

One key concept of inner model theory is building *iterated ultrapowers* to compare two models.




Comparison

One key concept of inner model theory is building *iterated ultrapowers* to compare two models.




Comparison

One key concept of inner model theory is building *iterated ultrapowers* to compare two models.




Comparison

One key concept of inner model theory is building *iterated ultrapowers* to compare two models.




Comparison

One key concept of inner model theory is building *iterated ultrapowers* to compare two models.




Comparison

One key concept of inner model theory is building *iterated ultrapowers* to compare two models.




Comparison

One key concept of inner model theory is building *iterated ultrapowers* to compare two models.



Comparison

One key concept of inner model theory is building *iterated ultrapowers* to compare two models.



Iteration trees

If the models contain Woodin cardinals, a linear iteration is not enough.

Iteration trees

If the models contain Woodin cardinals, a linear iteration is not enough. We construct a *tree of models* with embeddings along the branches by building ultrapowers at successor steps and direct limits along the branches at limit steps.


Iteration trees

If the models contain Woodin cardinals, a linear iteration is not enough. We construct a *tree of models* with embeddings along the branches by building ultrapowers at successor steps and direct limits along the branches at limit steps.

\mathcal{M}_0


Iteration trees

If the models contain Woodin cardinals, a linear iteration is not enough. We construct a *tree of models* with embeddings along the branches by building ultrapowers at successor steps and direct limits along the branches at limit steps.




Iteration trees

If the models contain Woodin cardinals, a linear iteration is not enough. We construct a *tree of models* with embeddings along the branches by building ultrapowers at successor steps and direct limits along the branches at limit steps.




Iteration trees

If the models contain Woodin cardinals, a linear iteration is not enough. We construct a *tree of models* with embeddings along the branches by building ultrapowers at successor steps and direct limits along the branches at limit steps.



Iteration trees

If the models contain Woodin cardinals, a linear iteration is not enough. We construct a *tree of models* with embeddings along the branches by building ultrapowers at successor steps and direct limits along the branches at limit steps.



The central problem is to choose a cofinal branch such that the direct limit is well-founded.

The iteration game

More precisely we consider the following two player game $\mathcal{G}(\mathcal{M}, \omega_1)$ of length $< \omega_1$ for a premouse \mathcal{M} .

The iteration game

More precisely we consider the following two player game $\mathcal{G}(\mathcal{M}, \omega_1)$ of length $< \omega_1$ for a premouse \mathcal{M} .

Player I plays at the successor steps: Choose an extender and apply it to the earliest model possible.

The iteration game

More precisely we consider the following two player game $\mathcal{G}(\mathcal{M}, \omega_1)$ of length $< \omega_1$ for a premouse \mathcal{M} .

- Player I** plays at the successor steps: Choose an extender and apply it to the earliest model possible.
- Player II** plays at the limit steps: Choose a cofinal well-founded branch of the iteration tree constructed so far.

The iteration game

More precisely we consider the following two player game $\mathcal{G}(\mathcal{M}, \omega_1)$ of length $< \omega_1$ for a premouse \mathcal{M} .

Player I plays at the successor steps: Choose an extender and apply it to the earliest model possible.

Player II plays at the limit steps: Choose a cofinal well-founded branch of the iteration tree constructed so far.

Then player II wins if all the models constructed in the game are well-founded.

The iteration game

More precisely we consider the following two player game $\mathcal{G}(\mathcal{M}, \omega_1)$ of length $< \omega_1$ for a premouse \mathcal{M} .

Player I plays at the successor steps: Choose an extender and apply it to the earliest model possible.

Player II plays at the limit steps: Choose a cofinal well-founded branch of the iteration tree constructed so far.

Then player II wins if all the models constructed in the game are well-founded.

Definition

We say a premouse \mathcal{M} is ω_1 -iterable iff player II has a winning strategy in the game $\mathcal{G}(\mathcal{M}, \omega_1)$. This winning strategy is called an iteration strategy for \mathcal{M} .



Inner Model Theory

Determinacy

Combinatorics of κ -
Trees

Looking beyond

Definition (Gale-Stewart 1953)

Let $A \subset \omega^\omega$. We denote the following game by $G(A)$

$$\begin{array}{c|cccc} \text{I} & n_0 & n_2 & \dots & \\ \hline \text{II} & & n_1 & n_3 & \dots \end{array} \quad \text{for } n_k \in \omega \text{ for all } k \in \omega.$$

We say player I wins the game iff $(n_k)_{k \in \omega} \in A$. Otherwise player II wins. We say $G(A)$ (or A itself) is *determined* iff one of the players has a winning strategy (in the obvious sense).

Which games are determined and what is it good for?

Theorem (Gale-Stewart, 1953)

(AC) Let $A \subset \omega^\omega$ be open or closed. Then $G(A)$ is determined.

Which games are determined and what is it good for?

Theorem (Gale-Stewart, 1953)

(AC) Let $A \subset \omega^\omega$ be open or closed. Then $G(A)$ is determined.

Theorem (Gale-Stewart, 1953)

Assuming AC there is a set of reals which is not determined.

Which games are determined and what is it good for?

Theorem (Gale-Stewart, 1953)

(AC) Let $A \subset \omega^\omega$ be open or closed. Then $G(A)$ is determined.

Theorem (Gale-Stewart, 1953)

Assuming AC there is a set of reals which is not determined.

Determinacy implies regularity properties.

Theorem (Mycielski, Swierczkowski, Mazur, Davis)

If all sets of reals are determined, then all sets of reals are Lebesgue measurable, have the Baire property, and have the perfect set property.

Determinacy for Definable Sets of Reals

Theorem (Martin, 1975)

Assume ZFC. Then every Borel set of reals is determined.

Determinacy for Definable Sets of Reals

Theorem (Martin, 1975)

Assume ZFC. Then every Borel set of reals is determined.

Determinacy for all analytic sets of reals is not provable in ZFC alone.

Theorem (Martin, 1970)

Assume ZFC and that there is a measurable cardinal. Then every analytic set is determined.

Determinacy for Definable Sets of Reals

Theorem (Martin, 1975)

Assume ZFC. Then every Borel set of reals is determined.

Determinacy for all analytic sets of reals is not provable in ZFC alone.

Theorem (Martin, 1970)

Assume ZFC and that there is a measurable cardinal. Then every analytic set is determined.

Theorem (Martin-Steel, 1985)

Assume ZFC and there are n Woodin cardinals with a measurable cardinal above them all. Then every Σ_{n+1}^1 set is determined.

Determinacy for Definable Sets of Reals

Theorem (Martin, 1975)

Assume ZFC. Then every Borel set of reals is determined.

Determinacy for all analytic sets of reals is not provable in ZFC alone.

Theorem (Martin, 1970)

Assume ZFC and that there is a measurable cardinal. Then every analytic set is determined.

Theorem (Martin-Steel, 1985)

Assume ZFC and there are n Woodin cardinals with a measurable cardinal above them all. Then every Σ_{n+1}^1 set is determined.

Are large cardinals necessary for the determinacy of these sets of reals?
How can these large cardinals affect what happens with the sets of reals?

Iterable Models and Projective Determinacy

Theorem (Neeman, Woodin)

Let $n \geq 1$. Then the following are equivalent.

- (a) Σ_{n+1}^1 -determinacy.
- (b) For every $x \in \mathbb{R}$ the ω_1 -iterable countable model of set theory with n Woodin cardinals $M_n^\#(x)$ exists.

Theorem (Neeman, Woodin)

Let $n \geq 1$. Then the following are equivalent.


- (a) Σ_{n+1}^1 -determinacy.
- (b) For every $x \in \mathbb{R}$ the ω_1 -iterable countable model of set theory with n Woodin cardinals $M_n^\#(x)$ exists.



Theorem (Neeman, Woodin)

Let $n \geq 1$. Then the following are equivalent.


- (a) Σ_{n+1}^1 -determinacy.
- (b) For every $x \in \mathbb{R}$ the ω_1 -iterable countable model of set theory with n Woodin cardinals $M_n^\#(x)$ exists.



Theorem (Neeman, Woodin)

Let $n \geq 1$. Then the following are equivalent.


- (a) Σ_{n+1}^1 -determinacy.
- (b) For every $x \in \mathbb{R}$ the ω_1 -iterable countable model of set theory with n Woodin cardinals $M_n^\#(x)$ exists.



Theorem (Neeman, Woodin)

Let $n \geq 1$. Then the following are equivalent.

- (a) Σ_{n+1}^1 -determinacy.
- (b) For every $x \in \mathbb{R}$ the ω_1 -iterable countable model of set theory with n Woodin cardinals $M_n^\#(x)$ exists.



For (a) \Rightarrow (b) see (M, Schindler, Woodin) “Mice with Finitely many Woodin Cardinals from Optimal Determinacy Hypotheses”, JML 2020.

For (b) \Rightarrow (a) see (Neeman) “Optimal proofs of determinacy II”, JML 2002.

Why stop playing at ω ?

Longer Games

Why stop playing at ω ? Define more generally:

Definition (Gale-Stewart 1953)

Let $A \subset \omega^\alpha$ for some ordinal α . We denote the following game by $G_\alpha(A)$

I	n_0	n_2	\dots	n_ω	\dots
II	n_1	n_3	\dots	$n_{\omega+1}$	\dots

for $n_\beta \in \omega$ for all $\beta < \alpha$.

As before, we say player I wins the game iff $(n_\beta)_{\beta < \alpha} \in A$. Otherwise player II wins. Moreover, $G_\alpha(A)$ (or A itself) is *determined* iff one of the players has a winning strategy (in the obvious sense).

Write $\text{Det}_\alpha(\Lambda)$ for the statement “all games of length α with payoff in Λ are determined”.

Theorem (Mycielski, 1964)

AD_{ω_1} , *determinacy for arbitrary games of length ω_1 , is inconsistent.*

Theorem (Mycielski, 1964)

AD_{ω_1} , *determinacy for arbitrary games of length ω_1 , is inconsistent.*

Proposition

$\text{Det}_{\omega \cdot (n+1)}(\mathbf{\Pi}_1^1)$ implies $\text{Det}_{\omega}(\mathbf{\Pi}_{n+1}^1)$.

Idea: “Simulate” projections by ω moves in a longer game, where we only consider the moves of one of the two players.

Theorem (Neeman, 2004)

Let $\alpha > 1$ be a countable ordinal and suppose that there are $-1 + \alpha$ Woodin cardinals with a measurable cardinal above them all. Then $\text{Det}_{\omega \cdot \alpha}(\mathbf{\Pi}_1^1)$ holds.

Theorem (Neeman, 2004)

Let $\alpha > 1$ be a countable ordinal and suppose that there are $-1 + \alpha$ Woodin cardinals with a measurable cardinal above them all. Then $\text{Det}_{\omega \cdot \alpha}(\mathbf{\Pi}_1^1)$ holds.

Is this result optimal?

Theorem (Neeman, 2004)

Let $\alpha > 1$ be a countable ordinal and suppose that there are $-1 + \alpha$ Woodin cardinals with a measurable cardinal above them all. Then $\text{Det}_{\omega \cdot \alpha}(\mathbf{\Pi}_1^1)$ holds.

Is this result optimal?

Let's focus on the first interesting level $\alpha = \omega + 1$.

Theorem (Aguilera-M, JSL 2020)

Suppose $\text{Det}_{\omega \cdot (\omega+1)}(\mathbf{\Pi}_1^1)$. Then there is a premouse with $\omega + 1$ Woodin cardinals.

In fact, the proof only uses $\text{Det}_{\omega^2}(\mathbf{\Pi}_2^1)$.

Theorem (Trang, 2013, building on Woodin)

Let α be a countable ordinal and suppose $\text{Det}_{\omega^{1+\alpha}}(\mathbf{\Pi}_1^1)$. Then there is a premouse with ω^α Woodin cardinals.

Larger Countable Ordinals

Theorem (Trang, 2013, building on Woodin)

Let α be a countable ordinal and suppose $\text{Det}_{\omega^{1+\alpha}}(\mathbf{\Pi}_1^1)$. Then there is a premouse with ω^α Woodin cardinals.

Theorem (M, 2019)

Let α be a countable ordinal and suppose $\text{Det}_{\omega^{1+\alpha}}(\mathbf{\Pi}_{n+1}^1)$. Then there is a premouse with $\omega^\alpha + n$ Woodin cardinals.

Larger Countable Ordinals

Theorem (Trang, 2013, building on Woodin)

Let α be a countable ordinal and suppose $\text{Det}_{\omega^{1+\alpha}}(\mathbf{\Pi}_1^1)$. Then there is a premouse with ω^α Woodin cardinals.

Theorem (M, 2019)

Let α be a countable ordinal and suppose $\text{Det}_{\omega^{1+\alpha}}(\mathbf{\Pi}_{n+1}^1)$. Then there is a premouse with $\omega^\alpha + n$ Woodin cardinals.

These methods should allow an analysis of the large cardinal strength of all analytic games of fixed countable length.



Inner Model Theory

Determinacy

Combinatorics of κ -
Trees

Looking beyond

This part is joint work with Yair Hayut.

Definition

Let κ be a regular cardinal. A tree T of height κ is called a *normal κ -tree* if

- each level of T has size $< \kappa$,
- every node splits,
- for every $t \in T$ and $\alpha < \kappa$ above the height of t , there is some t' of level α in T such that $t <_T t'$, and
- for every limit ordinal $\alpha < \kappa$ and every branch up to α there is at most one least upper bound in T .

The Branch Spectrum

Definition

Let κ be a regular cardinal. The *Branch Spectrum* of κ is the set

$$\mathfrak{S}_\kappa = \{ |[T]| \mid T \text{ is a normal } \kappa\text{-tree} \}.$$

The Branch Spectrum

Definition

Let κ be a regular cardinal. The *Branch Spectrum* of κ is the set

$$\mathfrak{S}_\kappa = \{|[T]| \mid T \text{ is a normal } \kappa\text{-tree}\}.$$

Examples

- $\mathfrak{S}_\omega = \{2^{\aleph_0}\}.$

The Branch Spectrum

Definition

Let κ be a regular cardinal. The *Branch Spectrum* of κ is the set

$$\mathfrak{S}_\kappa = \{|T| \mid T \text{ is a normal } \kappa\text{-tree}\}.$$

Examples

- $\mathfrak{S}_\omega = \{2^{\aleph_0}\}$.
- For $\kappa > \omega$, there are no κ -Kurepa trees iff $\max(\mathfrak{S}_\kappa) = \kappa$.

The Branch Spectrum

Definition

Let κ be a regular cardinal. The *Branch Spectrum* of κ is the set

$$\mathfrak{S}_\kappa = \{|[T]| \mid T \text{ is a normal } \kappa\text{-tree}\}.$$

Examples

- $\mathfrak{S}_\omega = \{2^{\aleph_0}\}$.
- For $\kappa > \omega$, there are no κ -Kurepa trees iff $\max(\mathfrak{S}_\kappa) = \kappa$.
- For $\kappa > \aleph_1$ of uncountable cofinality, tree property holds at κ iff $\min(\mathfrak{S}_\kappa) = \kappa$.

Upper Bounds

Let $\kappa > \aleph_1$.

Branch Spectrum	Upper bound
$\max(\mathfrak{S}_\kappa) = \kappa$ or $\kappa^+ \notin \mathfrak{S}_\kappa$	inaccessible cardinal

Upper Bounds

Let $\kappa > \aleph_1$.

Branch Spectrum	Upper bound
$\max(\mathfrak{S}_\kappa) = \kappa$ or $\kappa^+ \notin \mathfrak{S}_\kappa$	inaccessible cardinal
$\min(\mathfrak{S}_\kappa) = \kappa$	weakly compact cardinal

Upper Bounds

Let $\kappa > \aleph_1$.

Branch Spectrum	Upper bound
$\max(\mathfrak{S}_\kappa) = \kappa$ or $\kappa^+ \notin \mathfrak{S}_\kappa$	inaccessible cardinal
$\min(\mathfrak{S}_\kappa) = \kappa$	weakly compact cardinal
$\kappa^+ \notin \mathfrak{S}_\kappa$ and $\min(\mathfrak{S}_\kappa) = \kappa$?

Upper Bounds

Let $\kappa > \aleph_1$.

Branch Spectrum	Upper bound
$\max(\mathfrak{S}_\kappa) = \kappa$ or $\kappa^+ \notin \mathfrak{S}_\kappa$	inaccessible cardinal
$\min(\mathfrak{S}_\kappa) = \kappa$	weakly compact cardinal
$\kappa^+ \notin \mathfrak{S}_\kappa$ and $\min(\mathfrak{S}_\kappa) = \kappa$?

The following gives an upper bound.

Proposition

Let κ be $<\mu$ -supercompact, where μ is strongly inaccessible. Then, there is a forcing extension in which κ is weakly compact, $\mathfrak{S}_\kappa = \{\kappa, \kappa^{++}\}$.

Proof idea: Consider $\text{Col}(\kappa, <\mu) \times \text{Add}(\kappa, \mu^+)$.

Upper Bounds

Let $\kappa > \aleph_1$.

Branch Spectrum	Upper bound
$\max(\mathfrak{S}_\kappa) = \kappa$ or $\kappa^+ \notin \mathfrak{S}_\kappa$	inaccessible cardinal
$\min(\mathfrak{S}_\kappa) = \kappa$	weakly compact cardinal
$\kappa^+ \notin \mathfrak{S}_\kappa$ and $\min(\mathfrak{S}_\kappa) = \kappa$?

The following gives an upper bound.

Proposition

Let κ be $<\mu$ -supercompact, where μ is strongly inaccessible. Then, there is a forcing extension in which κ is weakly compact, $\mathfrak{S}_\kappa = \{\kappa, \kappa^{++}\}$.

Proof idea: Consider $\text{Col}(\kappa, <\mu) \times \text{Add}(\kappa, \mu^+)$.

Question

Is this optimal?

A First Lower Bound

If for some weakly compact cardinal κ , $\kappa^+ \notin \mathfrak{S}_\kappa$ then $0^\#$ exists:

A First Lower Bound

If for some weakly compact cardinal κ , $\kappa^+ \notin \mathfrak{S}_\kappa$ then $0^\#$ exists:

Fact (essentially Solovay)

If $0^\#$ does not exist then every weakly compact cardinal carries a tree with κ^+ many branches.

Theorem (Hayut, M.)

Let κ be a weakly compact cardinal and let us assume that there is no κ -tree with exactly κ^+ many branches. Then there is a non-domestic mouse. In particular, there is a model of $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$.

Theorem (Hayut, M.)

Let κ be a weakly compact cardinal and let us assume that there is no κ -tree with exactly κ^+ many branches. Then there is a non-domestic mouse. In particular, there is a model of $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$.

Proof idea:

- Consider a tree $\mathbb{T}(\mathcal{S})$ in $\mathcal{S} = \mathcal{S}(\kappa)$ the stack of mice on $K^c \parallel \kappa$ (cf. Andretta-Neeman-Steel and Jensen-Schimmerling-Schindler-Steel).

Theorem (Hayut, M.)

Let κ be a weakly compact cardinal and let us assume that there is no κ -tree with exactly κ^+ many branches. Then there is a non-domestic mouse. In particular, there is a model of $\text{ZF} + \text{DC} + \text{AD}_{\mathbb{R}}$.

Proof idea:

- Consider a tree $\mathbb{T}(\mathcal{S})$ in $\mathcal{S} = \mathcal{S}(\kappa)$ the stack of mice on $K^c \parallel \kappa$ (cf. Andretta-Neeman-Steel and Jensen-Schimmerling-Schindler-Steel).
- $\mathbb{T}(\mathcal{S})$ has exactly $(\kappa^+)^V$ many branches (using covering as in JSSS).

What is the large cardinal strength of this statement?

Conjecture

Let κ be a weakly compact cardinal such that there is no κ -tree with exactly κ^+ many branches. Then there is an inner model with a pair of cardinals $\lambda < \mu$ such that λ is $<\mu$ -supercompact and μ is inaccessible.



Inner Model Theory

Determinacy

Combinatorics of κ -
Trees

Looking beyond

Where is the problem when trying to prove such conjectures?

Where is the problem when trying to prove such conjectures?

- We currently do not know how to construct an inner model with a supercompact cardinal.

Where is the problem when trying to prove such conjectures?

- We currently do not know how to construct an inner model with a supercompact cardinal.
- But even if we were able to do that, we currently do not know how to “reach it”.

What does that mean?

What does that mean?

- Most arguments in inner model theory work bottom up. E.g., we prove small lower bounds and then argue inductively. We have seen this in the argument for mice from projective determinacy.

What does that mean?


- Most arguments in inner model theory work bottom up. E.g., we prove small lower bounds and then argue inductively. We have seen this in the argument for mice from projective determinacy.
- This leads to the *Core Model Induction* (CMI) technique.

What does that mean?

- Most arguments in inner model theory work bottom up. E.g., we prove small lower bounds and then argue inductively. We have seen this in the argument for mice from projective determinacy.
- This leads to the *Core Model Induction* (CMI) technique.
- Higher levels of this technique build on the *triple helix* of descriptive inner model theory:

What does that mean?

- Most arguments in inner model theory work bottom up. E.g., we prove small lower bounds and then argue inductively. We have seen this in the argument for mice from projective determinacy.
- This leads to the *Core Model Induction* (CMI) technique.
- Higher levels of this technique build on the *triple helix* of descriptive inner model theory:



What does that mean?

- 1st problem (Sargsyan, Trang, 2019): The current techniques of the CMI cannot go past *Sealing*, an hypothesis weaker than a Woodin cardinal that is a limit of Woodin cardinals

What does that mean?

- 1st problem (Sargsyan, Trang, 2019): The current techniques of the CMI cannot go past *Sealing*, an hypothesis weaker than a Woodin cardinal that is a limit of Woodin cardinals
- 2nd problem: For large enough large cardinals (e.g., a Woodin cardinal that is a limit of Woodin cardinals) there is no corresponding determinacy hypothesis known in the triple helix.

What does that mean?

- 1st problem (Sargsyan, Trang, 2019): The current techniques of the CMI cannot go past *Sealing*, an hypothesis weaker than a Woodin cardinal that is a limit of Woodin cardinals
- 2nd problem: For large enough large cardinals (e.g., a Woodin cardinal that is a limit of Woodin cardinals) there is no corresponding determinacy hypothesis known in the triple helix.

→ The hierarchy of long games could help here.



Inner Model Theory

Determinacy

Combinatorics of κ -
Trees

Looking beyond



Inner Model Theory

Determinacy

Combinatorics of κ -
Trees

Looking beyond

“There is an ever changing list of questions in set theory the answers to which would greatly increase our understanding of the universe of sets. The difficulty of course is the ubiquity of independence: almost always the questions are independent.”

(W. H. Woodin in Suitable Extender Models I)